

# Spatial depth for object-valued data

**J. Virta**

University of Turku

11th Tartu Conference on Multivariate Statistics, 26th of June  
2024

# Reference

This talk is based on the following preprint:

- **Virta, Joni.** *Spatial depth for data in metric spaces.* arXiv preprint arXiv:2306.09740 (2023).

# Table of Contents

- 1 Spatial depth
- 2 Object-valued data
- 3 Spatial depth for object-valued data
- 4 Data example
- 5 Closing remarks

# Ordering in $\mathbb{R}$


- Univariate samples of points admit a natural ordering.



- This ordering lets us identify **central or outlying points**.

# Ordering in $\mathbb{R}^2$

- How can we order a bivariate sample?



# Spatial depth

- Spatial depth [Chaudhuri, 1996, Vardi and Zhang, 2000] offers a way to order points in  $\mathbb{R}^P$ .
- For a point  $\mu \in \mathbb{R}^P$  and a random vector  $X \sim P$ , we define


$$D(\mu; P) := 1 - \left\| \mathbb{E} \left( \frac{X - \mu}{\|X - \mu\|} \right) \right\|^2.$$

- Intuitively,


$$\left\| \mathbb{E} \left( \frac{X - \mu}{\|X - \mu\|} \right) \right\|$$

is the **length of the average of unit-length arrows** drawn from  $\mu$  towards the sample points.

# Example




# Example






# Example



# Spatial depths of all points

The points are divided into **central** and **outlying**.



# Table of Contents

- 1 Spatial depth
- 2 Object-valued data**
- 3 Spatial depth for object-valued data
- 4 Data example
- 5 Closing remarks

# Object-valued data

- Nowadays, many samples  $X_1, \dots, X_n$  can be seen to consist of **objects**, such as
  - images,
  - functions,
  - graphs,
  - correlation matrices,
  - ...
- Often the **difference between two objects** is easy to quantify.

# Analysing object data

- A natural mathematical framework for object-valued data:
  - ① A metric space  $(\mathcal{X}, d)$ .
  - ② A random variable  $X \sim P$  taking values in  $\mathcal{X}$ .
  - ③ Methodology that relies on  $X$  only through the metric  $d(\cdot, \cdot)$ .
- Object methods are **extremely general**.
- Often, object methodology is constructed such that some familiar method is recovered when  $(\mathcal{X}, d)$  is Euclidean.

# Table of Contents

- 1 Spatial depth
- 2 Object-valued data
- 3 Spatial depth for object-valued data**
- 4 Data example
- 5 Closing remarks

# Metric spatial depth

- Let  $h : \mathcal{X}^3 \rightarrow \mathbb{R}$  be defined as

$$h(x_1, x_2, x_3) := \mathbb{I}(x_3 \notin \{x_1, x_2\}) \frac{d^2(x_1, x_3) + d^2(x_2, x_3) - d^2(x_1, x_2)}{d(x_1, x_3)d(x_2, x_3)},$$

where  $\mathbb{I}(\cdot)$  denotes the indicator function.

## Metric spatial depth [Virta, 2023]

$$D(\mu; P) := 1 - \frac{1}{2} \mathbb{E}\{h(X_1, X_2, \mu)\},$$

where  $X_1, X_2 \sim P$  are independent.

# Theoretical properties

## Property 1

When  $(\mathcal{X}, d)$  is an Euclidean space, then the metric spatial depth equals the classical spatial depth.



# Theoretical properties

## Property 2

$D(\mu; P)$  is finite for all  $\mu \in \mathcal{X}$  and all  $P$ .

## Property 3

The influence function of  $D$  satisfies

$$\sup_{z \in \mathcal{X}} |IF(z; D, \mu, P)| \leq 4.$$

# Theoretical properties

Let  $L[x, y, z]$  denote the event that

$$d(x, z) = d(x, y) + d(y, z),$$

i.e., that the point  $y$  is along the way from  $x$  to  $z$

## Property 4

Assume that  $P$  has no atoms. Then,

- (i)  $D(\mu; P) \in [0, 2]$ .
- (ii)  $D(\mu; P) = 0$  if and only if  $P(L[X_1, X_2, \mu] \cup L[X_2, X_1, \mu]) = 1$ .
- (iii)  $D(\mu; P) = 2$  if and only if  $P(L[X_1, \mu, X_2]) = 1$ .
- (iv) If  $(\mathcal{X}, d)$  is a Hilbert space, then  $D(\mu; P) \leq 1$ .

# Theoretical properties

## Property 5

Let  $\mu_n$  be a divergent sequence in  $\mathcal{X}$ . Then  $D(\mu_n; P) \rightarrow 0$  as  $n \rightarrow \infty$ .

# Theoretical properties

## Property 6

Assume that  $P$  has no atoms. Then  $D(\mu; P)$  is continuous in both  $\mu$  and  $P$ .





# Table of Contents

- 1 Spatial depth
- 2 Object-valued data
- 3 Spatial depth for object-valued data
- 4 Data example**
- 5 Closing remarks

# Non-convex depth regions

- It is not necessary to use the Euclidean metric with data in  $\mathbb{R}^p$  and we can instead use **metrics that promote non-linearity**

# Depth regions



# Table of Contents

- 1 Spatial depth
- 2 Object-valued data
- 3 Spatial depth for object-valued data
- 4 Data example
- 5 Closing remarks**






# Literature

- A review of the current state of object-valued data analysis can be found in:

**Dubey Paromita, Yaqing Chen, and Hans-Georg Müller.**  
*Metric statistics: Exploration and inference for random objects with distance profiles.* Annals of Statistics 52.2 (2024): 757-792.

Thank you for your attention!

# References I

-  Chaudhuri, P. (1996).  
On a geometric notion of quantiles for multivariate data.  
*Journal of the American Statistical Association*,  
91(434):862–872.
-  Li, J., Cuesta-Albertos, J. A., and Liu, R. Y. (2012).  
DD-classifier: Nonparametric classification procedure based on  
DD-plot.  
*Journal of the American Statistical Association*,  
107(498):737–753.
-  Vardi, Y. and Zhang, C.-H. (2000).  
The multivariate  $L_1$ -median and associated data depth.  
*Proceedings of the National Academy of Sciences*,  
97(4):1423–1426.

## References II



Virta, J. (2023).

Spatial depth for data in metric spaces.

*arXiv preprint arXiv:2306.09740.*



Xiao, H., Rasul, K., and Vollgraf, R. (2017).

Fashion-MNIST: a novel image dataset for benchmarking machine learning algorithms.

*arXiv preprint arXiv:1708.07747.*

# Table of Contents

6 Example metric spaces


7 Classification example

# Example I

- Assume that  $P$  puts equal mass  $1/n$  to each of the fixed points  $z_1, \dots, z_n \in \mathcal{X}$ .
- Question: Under what metric is the depth  $D(z_1; P)$  maximal?


# Example 1

- The railway metric uniquely gives  
$$D(z_1; P) = 1 + \left(1 - \frac{1}{n}\right) \left(1 - \frac{3}{n}\right).$$



# Example II

- Equip the finite set  $\mathcal{X} = \{1, \dots, n\}$  with the discrete metric  $d(i, j) = 1 - \mathbb{I}(i = j)$ .



- For which probability distribution  $P = (p_1, \dots, p_n)$  is  $D(1; P)$  maximized/minimized?



# Example II

- (i)  $D(1; P) = 1$  if and only if  $p_1 = 1$ .
- (ii)  $D(1; P) = 0$  if and only if exactly one of  $p_2, \dots, p_n$  equals 1.

# Table of Contents

6 Example metric spaces

7 Classification example

# Depth-depth plot

- Let  $P_{n1}$  and  $P_{n2}$  be the empirical distributions of two samples corresponding to different groups.
- In DD-classification [Li et al., 2012], we compute the depth vectors

$$z_i := (D(x_i; P_{n1}), D(x_i; P_{n2})).$$

- A test point  $x \in \mathcal{X}$  is then classified based on the vector

$$z := (D(x; P_{n1}), D(x; P_{n2})).$$

# Data example


- We applied DD-classification to a subsample of the FashionMNIST data set [Xiao et al., 2017] consisting of images of dresses, shirts and ankle boots.




# Specifications

- We randomly drew a training sample of  $n = 150$  images and a test sample of  $n_0 = 50$  images and used DD-classifier to predict the labels of the test images.
- We used metric spatial depth with the  $L_p$ -distance with  $p = 0.5, 0.6, \dots, 5$  as the metric.
- We considered both LDA and QDA.
- The experiment was repeated a total of 100 times.

# Results



# DD-plot



# Interpretation

- Both LDA and QDA reach their maximal performance at super-Euclidean geometry.
- QDA is uniformly superior to LDA.